



44

"Express Mail" mailing label number EL544993670US

PATENT
Case U 0069 CC/SPAP

COPY OF PAPERS
ORIGINALLY FILED

ASTRINGENT SHAVE PREPARATIONS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims the benefit of copending provisional application serial number 60/246,355 filed on November 7, 2000.

5 STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

Not applicable.

BACKGROUND OF THE INVENTION

10 The present invention generally relates to shaving preparations which are applied onto a person's skin prior to shaving. A common problem associated with the act of shaving is the formation of razor bumps on the shaved surface, i.e., pseudofolliculitis barbae (PFB). This is a shaving-induced condition caused by re-entrant penetration of newly-cut hairs into the skin resulting in significant
15 skin irritation.

Hairs which lie nearly parallel to the skin have a tendency of being cut at an acute angle, producing sharp points that can penetrate into the skin. Those individuals whose hair follicles have a strong natural curl are even more susceptible to this type of skin condition. The re-entrant hair induces an
20 antigenic, foreign body reaction, leading to the formation of papules or cysts that appear as raised bumps on the skin's surface. These raised bumps are further

more upright, which also facilitates a closer shave and causes the shaved hairs to have blunt rather than sharp ends.

While any suitable astringent compound may be employed such as, for example, witch hazel extract or zinc paraphenolsulfonate, a particularly effective
5 astringent additive is a mixture of butylene glycol and *Fomes officinalis* (mushroom extract) commercially available from Laboratories Serobiologiques under the tradename LARICYL® LS 8865.

The astringent compound may be employed in any conventional type of shaving composition. A particularly preferred shaving composition is a shaving
10 gel. This type of composition is typically in the form of an oil-in-water emulsion containing water and a water-soluble soap component and/or synthetic wetting agent.

The primary component of the shaving gel is water, preferably deionized or distilled water. The soap component is typically a water soluble salt of a fatty
15 acid. Examples thereof include, but are not limited to, triethanolamine soaps of stearic and palmitic acids, animal-derived soaps, vegetable-derived soaps, and mixtures thereof. Synthetic wetting agents, when used, are typically of the nonionic variety. Examples thereof include, but are not limited to, water-soluble alkoxylated fatty alcohols, water soluble polyoxyethylene ethers of alkyl-
20 substituted phenols, and the like.

The shaving gel may also include a self-foaming agent, if desired, which provides for the formation of foam when spread over a person's skin and beard. Typically, a low boiling aliphatic hydrocarbon is used as the volatile self-foaming
agent. Examples thereof include, but are not limited to, n-pentane, isopentane,
25 neopentane, n-butane, isobutane, and mixtures thereof.

According to one aspect of the present invention, there is provided a shaving composition containing an astringent additive for use prior to shaving. The astringent additive may be present in the shaving composition in an amount of from about 0.1 to about 35% by weight, preferably from about 1 to about 20%

by weight, and most preferably from about 2 to about 10% by weight, all weights being based on the weight of the composition.

According to another aspect of the present invention, there is provided a process for inhibiting the re-entrant penetration of newly shaved hair into human skin. The process involves contacting the area to be shaved, prior to shaving, with the above-described shaving composition.

The present invention will be better understood by the following examples, all of which are intended to be illustrative only, and are not meant to unduly limit the scope of the invention.

EXAMPLES

Example 1

A non-foaming shaving gel was prepared per the following formulation.

<u>Component</u>	<u>% by weight</u>
water	91.04
ELESTAB® 338	2.50
hydroxyethyl cellulose	0.50
CARBOMER®	0.16
sodium hydroxide (1N)	0.80
LARICYL® LS 8865	5.00

The non-foaming shaving gel was applied onto human skin, prior to shaving, producing a strong tingling/tightening sensation during the contact time required for shaving.

Example 2

A foaming shave gel was prepared per the following formulation.

	<u>Component</u>	<u>% by weight</u>
	water	37.10
	ELESTAB® 338	1.00
	hydroxypropyl methylcellulose	0.60
5	triethanolamine (99%)	0.30
	cocamide MEA	5.50
	citric acid (50%)	0.30
	ammonium lauryl sulfate	28.00
	ammonium laureth sulfate	21.00
10	dimethicone copolyol	0.20
	lauryl glycoside	2.00
	cocoglycoside + glycerol oleate	1.00
	LARICYL® LS 8865	3.00

15 Example 3

A foaming shave gel was prepared per the following formulation.

	<u>Component</u>	<u>% by weight</u>
	water	35.80
20	ELESTAB® 338	1.00
	hydroxypropyl methylcellulose	0.50
	triethanolamine (99%)	0.30
	cocamide MEA	6.00
	ammonium lauryl sulfate	28.00
25	ammonium laureth sulfate	23.00
	dimethicone copolyol	0.20
	cocamidopropylamine oxide	2.00
	LARICYL® LS 8865	3.00